and suspected Lyme borreliosis. Neurology Sept 1997;49:817-824). (Reprints: Dr David Nadal, University Children's Hospital, Infectious Diseases Unit, Steinwiesstr. 75, CH-8032 Zurich, Switzerland).

COMMENT. Analysis of CSF for intrathecal specific antibodies to *B. burgdorferi* in children presenting with acute peripheral facial palsy may facilitate early diagnosis and prompt antibiotic treatment. In patients with lymphocytic pleocytosis but no intrathecal antibodies, detection of seroconversion by analysis of convalescent serum is important in diagnosis.

Neurologic signs and syndromes associated with Lyme disease in children are reviewed in three reports and commentaries in Progress in Pediatric Neurology III, PNB Publishers, 1997;pp484-5. In one series of 96 patients, facial palsy occurred in 14%. Pseudotumor, reviewed in the following report, was diagnosed in 6% of cases of Lyme disease.

PSEUDOTUMOR CEREBRI

INCIDENCE RATES OF PEDIATRIC PSEUDOTUMOR CEREBRI

Twenty nine cases of pseudotumor cerebri were identified in a retrospective study of records of 205,765 children aged 2-15 years presenting between 1979 and 1994 at the IWK Grace Health Centre, Dalhousie University, Halifax, Canada. The annual incidence of pseudotumor for the total group was 0.9 per 100,000 children; in age group 2-11 years, the incidence was 0.7, and for 12-15 years, 1.5. Females were affected nearly 3 times more frequently than males. The adolescent age group was affected twice as frequently as younger children. (Gordon N. Pediatric pseudotumor cerebri: descriptive epidemiology. Can J Neurol Sci Aug 1997;24:219-221). (Reprints: Dr Kevin Gordon, IWK-Grace Health Centre, 5850 University Ave, Box 3070, Halifax, Nova Scotia, Canada B3J 3G9).

COMMENT. The increasing incidence of pseudotumor cerebri in later childhood and adolescence (1.5 per 100,000), noted in this study of patients admitted after 1978, was also noteworthy in earlier reports. A review of the literature before 1960 showed that 84 (37%) of 224 patients of all ages were in children and of these 75 (90%) were between 5 and 15 years of age. (Millichap JG. Pediatrics Feb 1959;23:257). Antecedent otitis media was reported in 29%, and mild head injury or infection other than otitis occurred in an equal proportion of the cases. The advent of antibiotics virtually abolished the cases of "otitic hydrocephalus," and explained the increase in "idiopathic" cases in later reports. (See Progress in Pediatric Neurology III, PNB Publ, 1997;pp431-433, for articles on signs and symptoms, the MRI, and treatment of pseudotumor cerebri).

TRAUMATIC DISORDERS

POST-TRAUMATIC OLFACTORY DYSFUNCTION

The degree of olfactory function and damage to olfactory bulbs and tracts were quantitatively determined in 268 patients with head trauma (HT) presenting with complaints of olfactory disturbance at the University of Pennsylvania Smell and Taste Center, Philadelphia, PA. Anosmia occurred in 179 patients (67%), microsmia (lessened sense of smell) in 20%, and 12% had normal sense of smell. Olfactory test scores after head trauma were not related

to age or sex. Occipital or temporo-parietal impacts caused more cases of olfactory dysfunction than frontal impacts. Of 66 retested after periods ranging from 1 month to 13 years, 24 (36%) showed slight improvement, 45% were unchanged, and 18% were worse. Parosmia, or perversion of smell, occurring in 40% of patients immediately after the trauma, decreased to 15% during an 8 year follow-up. Volumes of olfactory bulbs and tracts measured by MRI were smaller than controls in male, but not female, patients. (Doty RL, Yousem DM, Pham LT et al. Olfactory dysfunction in patients with head trauma. Arch Neurol Sept 1997;54:1131-1140). (Reprints: Richard L Doty PhD, Smell and Taste Center, University of Pennsylvania Medical Center, 5 Ravdin Pavilion, 3400 Spruce St, Philadelphia, PA 19104).

COMMENT. Patients with head trauma who lose their sense of smell rarely regain normal olfactory function, whereas those who complain of distortions of smell usually recover over time. Head trauma male patients with olfactory dysfunction have MRI evidence of greatly reduced olfactory bulb and tract volumes. The apparent selective sparing of female olfactory structures may be explained by lesser severity of trauma or a protective effect of estrogens.

SEIZURE DISORDERS

X-LINKED INFANTILE SPASMS

Two unrelated families with X-linked infantile spasm syndrome were studied genetically by two-point and multipoint linkage analyses at the University Hospital Gasthuisberg, and Center for Human Genetics, University of Leuven, and University of Antwerp, Belgium. The disease gene was located to the distal part of the short arm of the X chromosome, between Xpter and Xp11.4. (Claes S, Devriendt K Lagae L et al. The X-linked infantile spasms syndrome (MIM 308350) maps to Xp11.4-Xpter in two pedigrees. Ann Neurol Sept 1997;42:360-364). (Respond: Dr JJ Cassiman, Center for Human Genetics, University of Leuven, Campus Gasthuisberg, Herestraat 49, B-3000 Leuven, Belgium).

COMMENT. West syndrome may rarely occur in families as an X-linked inherited disorder.

Zonisamide (ZNS) for infantile spasms. ZNS monotherapy was effective in the control of spasms in 4 of 11 patients treated at 11 centers in Japan. (Suzuki Y, Nagai T, Ono J et al. Epilepsia Sept 1997;38:1035-1038).

AUTOMATIC NEONATAL SEIZURE DETECTION BY EEG

EEGs obtained from 55 newborns, recorded at the Montreal, Sydney, and Texas Children's Hospitals, were reviewed by 3 types of automatic analysis of sequential epochs aimed at detecting rhythmic paroxysmal discharges, repetitive spike patterns, arrhythmic runs of spikes, and low frequency discharges, and the methods are reported from the Montreal Neurological Institute, and the Montreal Children's Hospital, Canada. Initial evaluation detected 71% of seizures and 78% of seizure clusters, and the false detection rate was 1.7%/hour of recording. (Gotman J, Flanagan D, Zhang J, Rosenblatt B. Automatic seizure detection in the newborn: methods and initial evaluation. Electroenceph clin Neurophysiol Sept 1997;103:356-362). (Respond: Dr J Gotman, Montreal Neurological Institute and Hospital, 3801 University St, Montreal, PQ, H3A 2B4 Canada).