PEDIATRIC NEUROLOGY BRIEFS

A MONTHLY JOURNAL REVIEW

J. GORDON MILLICHAP, M.D., F.R.C.P., EDITOR

Vol. 2, No. 4

April 1988

SEIZURE DISORDERS

EPIDEMIOLOGY OF EPILEPSY

All children 3 years of age on Jan 1, 1975 in Fuchu City, Tokyo, a total of 17,044 of a population of 182,000, were examined neurologically at the Health Center and the Dept of Genetics, Tokyo Metropolitan Institute for Neurosciences, Tokyo, Japan. Those with a history of seizures were compared with a randomly selected group of healthy children from the same population. The incidence of epilepsy (recurrent nonfebrile seizures) up to age 3 yrs was .43% and that of a single nonfebrile seizure was .47%. Febrile convulsions had occurred in 8.2%. By age 9-14 yrs the total cumulative incidence of epilepsy was .82%. Epilepsy developed in 50% of children with single nonfebrile seizures, in 2% of those with febrile convulsions, .2% of healthy 3-yr-old children, and in 2% of potential seizure patients with spike EEG abnormalities observed at 3 yrs of age. The author stresses identification and treatment of patients with potential seizure disorders. (Tsuboi T. Prevalence and incidence of epilepsy in Tokyo. Epilepsia March/April 1988; 29: 103-110).

COMMENT. This study provides detailed statistics of the incidence of febrile convulsions, nonfebrile seizures and epilepsy in Japan. Of particular interest are the relatively high frequency of febrile convulsions in Japan c.f. US and Europe (8.2 v 3%), and the low incidence of epilepsy among potential healthy carriers with EEG seizure activity in the general population (2%). Methods for the prevention of clinical seizures in potentially epileptic subjects were not addressed although this goal was proposed. Most authorities would not advocate the use of long-term prophylactic anticonvulsant medication in healthy children with abnormal EEGs or in those with a history of one febrile convulsion uncomplicated by nonfebrile seizures.

Two epidemiological studies of epilepsy in India reported in the same issue of Epilepsia showed prevalence rates of .3-.4% for

PEDIATRIC NEUROLOGY BRIEFS *1988 covers selected articles from the world literature and is published monthly. Subscription requests (\$25 US or £15 UK) may be sent to: Pediatric Neurology Briefs - J. Gordon Millichap, MD, FRCP - Editor, P.O. Box 11391, Chicago, IL 60611, USA, or P.O. Box 1605, London W8 7JD, UK. The Editor is Professor of Neurology and Pediatrics at Northwestern University Medical School, Chicago, Illinois, USA.

children and adolescent age groups. Males predominated and mental retardation and cerebral palsy were the most frequently associated conditions.

VARIABILITY OF SEIZURE CLASSIFICATION

One senior neurologist (JBB) from the Department of Neurology and Pediatrics, West Virginia University, Morgantown, W.Va., USA, and three neurology residents from the University of Oklahoma, Harvard University, and W.Va. University reviewed descriptions of seizures transcribed from medical records and classified them according to the International League Against Epilepsy (ILAE) Classification system, 1981. The overall agreement in classifying seizure types by the ILAE system was relatively poor and only slightly better than would have been expected by chance. Agreement was particularly weak for atypical absence, partial seizures with secondary generalization, and generalized motor seizures. It was better for simple and complex partial, simple absence and infantile spasms. Approximately 22% of descriptions were insufficient for classification by the ILAE system. The authors conclude that the development and testing of more explicit criteria for the diagnosis of specific seizure types may be especially useful in improving the reliability of seizure classification. (Bodensteiner JB et al. Interobserver variability in the ILAE classification of seizures in childhood. Epilepsia March/April 1988; 29: 123-128).

COMMENT. A classification of seizure types is of little value without assessment of its reproducibility in practice. Surprisingly, this study appears to be the first test of the reliability of the ILAE system of seizure classification, first proposed in 1970 and generally employed in statistical trials of anticonvulsant drugs. The relatively low level of concordance among observers using the ILAE system in the present study is disturbing, and the substitution of a more specific seizure classification should be entertained especially in therapeutic trials.

PSYCHOLOGICAL TREATMENT OF EPILEPSY

Three children, ages 12-15, with frequent refractory seizures were treated by behavioral intervention techniques in the Depts of Medical Rehabilitation and Clinical Neurophysiology, Orebro Medical Center Hospital, Orebro, and Dept of Clinical Psychology, University of Uppsala, Sweden. Seizure types described as myoclonic in pt 1, Jacksonian (pt 2), and minor motor (pt 3) were associated with mental deterioration. Behavioral countermeasures, consisting of tensing of muscle groups and screaming "stop"! (pt 1), massaging the affected limb (pt 2), and fixing eye movements on an object (pt 3), resulted in an immediate reduction in both the seizures and paroxysmal EEG activity in all three cases. Other behavioral strategies, including biofeedback training in awareness of early seizure signals, applied relaxation, and positive reinforcement, were without beneficial effect. The authors propose that countermeasures in the level and speed of arousal may be the mechanism of the treatment intervention. (Dahl J, Melin L, Leissner P. Effects of behavioral intervention on epileptic seizure behavior and paroxysmal activity: a systematic replication of three cases of children with intractable epilepsy. Epilepsia March/April 1988; 29: 172-183).