of symptomatic. A mean developmental score of 71 in cryptogenic cases was significantly higher than 48 in symptomatic. Additional forms of seizures were more common in symptomatic cases (57% cf 35%). Unfavorable prognostic factors, particularly cognitive deficits, were the association of other seizures, neurologic deficits, delay in treatment, poor response to ACTH, and persistent EEG abnormalities after 2 weeks of therapy. All patients in the cryptogenic group responded to ACTH (80 IU/IM/d for 2 weeks, tapered over 6 to 8 weeks). Of 30 symptomatic cases receiving ACTH, 12 (40%) were completely controlled initially, and 11 had a partial response. Response was higher in patients with developmental scores >70, both cryptogenic and symptomatic cases. (Koo B et al. Infantile spasms: Outcome and prognostic factors of cryptogenic and symptomatic groups. Neurology Nov 1993;43:2322-2327). (Reprints: Dr Betty Koo, Division of Neurology, The Hospital for Sick Children, 555 University Avenue, Toronto, ON, Canada M5G 1X8).

COMMENT. With improved diagnostic techniques, particularly the MRI, the increased recognition of cerebral dysgenesis as a frequent cause for infantile spasms has resulted in a lowered percentage of cryptogenic cases compared to earlier reports. In a study involving 61 cases at the Mayo Clinic (Millichap JG, Bickford RG, Klass DW, Backus RE. Epilepsia 1962;3:188-197), 43% were of undetermined etiology and only 7% were recognized with cerebral dysgenesis. The anticonvulsant efficacy and outcome of treatment with ACTH in 21 patients was related significantly to the age at the time of diagnosis and treatment; 80% of infants less than 1 year old were benefited whereas only 22% of those over 1 year showed reduction in seizures and EEG improvement. Control of seizures and hypsarrhythmia was not significantly related to the etiology, but patients of borderline or normal intelligence tended to respond more frequently than did those with developmental quotients <70. (Millichap JG, Bickford RG. JAMA 1962;182:523-527).

Posterior fossa abnormalities in children with infantile spasms were associated with a poor developmental outcome in a study of 98 children from New England Medical Center and Children's Hospital, Boston, and Hadassah University Hospital, Jerusalem (Schiffmann R et al. J Child Neurol Oct 1993;8:360-365). Brain stem dysfunction and atrophy were remarkable in a study of 10 patients with infantile spasms, employing evoked potentials and MRI, at the University of Tokushima, Japan (Miyazaki M et al. Neuropediatr June 1993;24:126-130).

BENIGN NEONATAL FAMILIAL CONVULSIONS

An electroencephalographic-video study of 14 benign neonatal familial convulsions (BNFCs) in 3 children from two families is reported from the Hopitaux Universitaires, Strasbourg, and the Institut de Biologie, Montpellier,

France. Seizures occurred during sleep and were preceded by a short period of arousal. A brief symmetrical flattening of the EEG, associated with apnea, tachycardia, and tonic motor activity, was followed by a long (1-2 minute) bilateral discharge of spikes and sharp waves, simultaneous with vocalizations, chewing, and focal or generalized clonic activity, often asymmetrical and asynchronous. EEG and clinical symptoms varied from left to right from one seizure to the next, and a stable focus was never observed. (Hirsch E et al. Electroclinical signs of benign neonatal familial convulsions. Ann Neurol Dec 1993;34:835-841). (Respond: Dr Hirsch, Service d'Epileptologie Clinique, Clinique Neurologique, Hopitaux Universitaires de Strasbourg, 1 place de l'Hopital, 67091 Strasbourg Cedex, France).

COMMENT. BNFCs are brief seizures beginning at the second or third day after birth and occurring for several days or weeks. Less than 10% have seizures in later childhood and adulthood. The gene for this syndrome has been localized to the long arm of chromosome 20, but BNFCs may be genetically heterogeneous. The pattern of the seizures is similar to that of generalized tonic-clonic epilepsy, in which the initial tonic phase may be asymmetrical. The asymmetry of BNFCs may be explained by immaturity of the corpus callosum in the neonate. The interictal EEG is either normal or shows bursts of sharp alternating theta rhythms, a theta pointu alternant pattern. Frequent BNFCs have no postictal deficit or exhaustion and no adverse effect on development of the child.

GENETICS OF UNVERRICHT-LUNDBORG MYOCLONUS EPILEPSY

Clinical and molecular genetic data on a family from the United States in which four of five sibs were affected with progressive myoclonus epilepsy of Unverricht-Lundborg type are reported from the University of Helsinki, Finland; National Cancer Institute, Bethesda, MD; and the VA Hospital, University of Florida, Gainesville, FL. Onset was between 9 and 11 years with grand mal seizures. Progressive stimulus-sensitive myoclonus developed at 11 to 13 years of age. Generalized spike-wave and polyspike-wave paroxysms and photo-sensitivity were characteristic EEG findings. The gene (EPM1) has been mapped to chromosome 21 in Finland, and the gene in this non-Finnish American family was also linked to the EPM1 region. Crossover events refined the gene localization between loci CBS and D21S112, leading to greater accuracy of genetic prediction based on linkage analysis. (Lehesjoki A-E et al. Progressive myoclonus epilepsy of Unverricht-Lundborg type: A clinical and molecular genetic study of a family from the United States with four affected sibs. Neurology Nov 1993;43:2384-2386). (Reprints: Dr Anna-Elina Lehesjoki, Dept Med Genetics, PO Box 21 (Haartmaninkatu 3), 00014 University of Helsinki, Finland).